

MA 181202

Roll No. of candidate

29/5/24

--	--	--	--	--	--	--	--	--	--

Bina Chowdhury Central Library
Girijananda Chowdhury University
2024 Matkhowapara, Azara, Ghy-17

B.Tech. 2nd Semester End-Term Examination

MATHEMATICS

**New Regulation (w.e.f. 2017-18) &
New Syllabus (w.e.f. 2018-19)**

Full Marks – 70

Time – Three hours

The figures in the margin indicate full marks
for the questions.

Answer question No. 1 and any *four* from the rest.

1. Select the correct answer :

(10 × 1 = 10)

(i) The divergence of $(3x^2i + 5xy^2j + xyz^3k)$ at the point (1, 2, 3) is

- (a) 20 (b) 40
(c) 80 (d) None of these

(ii) The unit normal vector to the surface $x^2y + 2xz = 4$ at the point (2, -2, 3) is

- (a) $\frac{1}{3}(-i + 2j + 2k)$ (b) $\frac{1}{3}(i - 2j + 2k)$
(c) $\frac{1}{3}(i + 2j - 2k)$ (d) none of these

(iii) The general solution of the equation $\frac{d^2y}{dx^2} + 3\frac{dy}{dx} + 2y = 0$ is

- (a) $y = Ae^{-x} + Be^{-2x}$ (b) $y = Ae^x + Be^{2x}$
(c) $y = e^{-x} + e^{-2x}$ (d) $y = (A + Bx)e^x$

(iv) The integrating factor of the differential equation $(1 + y^2)dx = (\tan^{-1}y - x)dy$ is

- (a) $e^{\tan^{-1}x}$ (b) $e^{\tan^{-1}y}$
(c) $\tan^{-1}x$ (d) $\tan^{-1}y$

[Turn over

(v) For the differential equation $(x^2 - 2x)\frac{d^2y}{dx^2} + \frac{dy}{dx} - 2y = 0$, the point $x = 0$ is

- (a) ordinary (b) regular singular
(c) irregular singular (d) none of these

(vi) For the Legendre polynomial $P_n(x)$, $P_n(-1)$ is

- (a) 1 (b) -1
(c) $(-1)^n$ (d) none of these

(vii) The real part of $f(z) = \cos z$ is

- (a) $\cosh x \cos y$ (b) $\cos x \cosh y$
(c) $\cos x \cosh x$ (d) none of these

(viii) For the function $f(z) = \frac{\cos z}{z}$, $z = 0$ is a

- (a) pole
(b) removable singularity
(c) essential singularity
(d) none of these

Bina Chowdhury Central Library
Girijananda Chowdhury University
Hatkowapara, Azara, Ghy-17

(ix) The image of the circle $|z| = 2$ under the mapping $w = z + (3 + 2i)$ is a

- (a) circle (b) ellipse
(c) pair of lines (d) hyperbola

(x) Which of the following functions is analytic?

- (a) $z + \bar{z}$ (b) $2z^2 + 2\bar{z} + c$
(c) $|z|^2$ (d) $z^2 + 2z + c$

2. (a) Find the directional derivative of the function $\phi = xy^2 + yz^3$ at the point $(2, -1, 1)$ in the direction of the normal to the surface $x \log z - y^2 + 4 = 0$ at $(-1, z, 1)$. (5)

(b) With the help of Green's theorem evaluate the line integral $\int_C \vec{A} \cdot d\vec{c}$ where

$\vec{A} = (x^2 + xy^2)\vec{i} + (y^2 + x^2y)\vec{j}$ and C is the boundary of the region bounded by $y = x$ and $y = x^2$. (5)

(c) Evaluate the surface integral of $\vec{A} = z\vec{i} + x\vec{j} - 3y^2z\vec{k}$ over the surface of the cylinder $x^2 + y^2 = 16$ included in the first octant between $z = 0$ and $z = 5$. (5)

3. (a) Solve the differential equation $(D-2)^2 y = 8(e^{2x} + \sin 2x)$. (6)

(b) Solve any THREE from the following : $(3 \times 3 = 9)$

(i) $(x^4 + y^4)dx - xy^3dy = 0$

(ii) $P = \sin(y - px)$

(iii) $(1+x^2)\frac{dy}{dx} + y = e \tan^{-1} x$

(iv) $y = 2px + p^4 x^2$.

Bina Chowdhury Central Library
Girijananda Chowdhury University
Hatkhowapara, Azara, Gny-17

4. (a) Solve in series the differential equation $\frac{d^2 y}{dx^2} + xy = 0$. (5)

(b) Express the polynomial $x^4 + x^3 + x^2 + 1$ in terms of Legendre polynomials. (5)

(c) Show that $J_{5/2}(x) = \sqrt{\frac{2}{\pi x}} \left(\frac{3-x^2}{x^2} \sin x - \frac{3}{x} \cos x \right)$. (5)

5. (a) Prove that $u = y^3 - 3x^2 y$ is a harmonic function. Construct its harmonic conjugate and find the corresponding analytic function $f(z)$ in terms of z . (5)

(b) Evaluate $\int_L \bar{z} dz$ from $z=0$ to $z=4i$, first along the straight line $z=0$ to $z=2i$ and then along the line to $z=4+2i$. (5)

(c) State Cauchy's integral formula and hence evaluate $\oint_C \frac{e^{-2z}}{(z+1)^4}$ where C is the circle $|z|=2$. (5)

6. (a) Consider the transformation $w = ze^{i\pi/4}$ and find the region in the w -plane corresponding to the triangular region bounded by the lines $x=0$, $y=0$ and $x+y=1$ in the z -plane. (5)

(b) Expand the function $f(z) = \frac{(z-2)(z+2)}{(z+1)(z+4)}$ in the region $1 < |z| < 4$. (5)

(c) State residue-theorem and apply it to evaluate $\oint_C \frac{z^2 dz}{(z-1)^2(z+2)}$ where C in the circle $|z|=3$. (5)

7. (a) State the generating function for Legendre polynomials and hence show that $P_n(-x) = (-1)^n P_n(x)$. (5)

(b) Use divergence theorem to show that $\oint_S V r^2 ds = 6V$ where S is any closed surface enclosing a volume V . (5)

(c) Evaluate $\int_0^{2\pi} \frac{d\theta}{5 + 4 \cos \theta}$. (5)

Bina Chowdhury Central Library
Girijananda Chowdhury University
Hatkhowapara, Azara, Gny-17